



nextmune

artuvet alergovet dr baddaky

Next+
Food Reactivity
Test

—




Food allergies

Diagnosis and management

Introduction

Adverse food reactions (AFR) encompass food intolerances as well as immunologically mediated hypersensitivities, that is, food allergies.



Pathogenesis

The specific immunological mechanisms involved in food allergy are not clearly understood. It is hypothesized that immunological food reactions (food allergy) develop when a food specific IgE antibody on a mast cell binds with a food antigen.

This binding initiates mast cell release of potent inflammatory mediators and cytokines¹. Although, it is believed that most food allergies are type I or IgE mediated, there is also some evidence to suggest Type III and IV hypersensitivity reactions may be involved².



nextmune



Prevalence, age of onset and genetic predisposition

Among animals with pruritus and clinical signs of allergic dermatitis, **the prevalence of AFR is high enough to justify this syndrome to be ruled-out**

The true incidence of such reactions is still not entirely clear, but has been reported to range between 10% and 25%, in dogs with allergic skin disease³. It is also estimated to be around 1/3 of dogs with atopic dermatitis³. It is described a stronger incidence in dogs under 1 year of age or above 6 years old⁴.

Labrador retrievers, West Highland White Terriers, Boxers, Rhodesian ridgebacks and Pug breeds are predisposed to developing AFR. The most common allergens responsible for AFR in dogs seemed to be beef, chicken, egg, milk, wheat, soy and corn, which are also common ingredients in many commercial foods⁵.



Clinical Signs

Dogs suffering from AFR generally present with dermatological or gastrointestinal signs, or combination of both.

Dermatological signs are varied, and often indistinguishable from those associated with atopic dermatitis, although the presence of an unusual distribution or concomitant gastrointestinal signs may raise suspicion for an adverse food reaction.

When pruritus is not corticosteroid-responsive, a food allergy should be considered (if pruritus responds to corticosteroids that does not rule out the possibility that a food allergy is present).

Generalized or localized non-seasonal pruritus (face, ears, paws, axillae, inguinal and perineal regions) is the most frequently described dermatological sign. It is also possible for the effects of a food allergy to be below the 'itch threshold' and only observe flares of



- Non-seasonal pruritus
- Signs of allergic dermatitis
- Gastrointestinal signs
- Corticosteroid non-responsive

pruritus with the addition of environmental allergens during high pollen season. Otitis externa and recurrent pyoderma with or without pruritus has also been associated with AFR in dogs.

Gastrointestinal signs may include vomiting, diarrhea, weight loss, and abdominal discomfort.

Ears

Appearance:

Waxy discharge; redness; odor

Behaviour:

Scratching or rubbing ears with paws or against furniture

Face

Appearance:

Inflammation; redness; conjunctivitis; swelling

Behaviour:

Scratching or rubbing face with paws or against furniture

Skin

Appearance:

Reddened; possibly with crusts or scales; odor with secondary infections

Behaviour:

Scratching and licking sides/belly, elbows, groin

- + Otitis externa
- Recurrent pyoderma
- Vomiting
- Diarrhea
- Weight loss
- Abdominal discomfort
- Flatulence

Feet

Appearance:

Inflammation, redness, odor, brown discoloration where licking has occurred

Behaviour:

Licking and chewing of feet/pads


Haircoat

Appearance:

Bald spots; brown discoloration where licking has occurred

Behaviour:

Scratching sides/belly, rubbing face against furniture or carpet, licking



Diagnosis

Currently, the definitive diagnosis of food allergy in dogs is only possible through a food elimination trial followed by a food challenge test. The recommended length of this trial is, at least, 6 weeks.

Complete or partial improvement of pruritus during the trial is the main criterion for suspected food allergy. The diagnosis is confirmed by the recurrence of clinical signs when the food given previously is reintroduced. As animals may be multi sensitive, it may be necessary to attempt dietary restriction more than once.

The elimination trial can be done with a homemade diet, a commercial diet with hydrolyzed protein, or a commercial diet with proteins unusual for the dog.


Commercial diets often contain larger protein molecules capable of inducing clinical deterioration in some dogs^{6,7} or protein sources not declared on the label^{8,9}.

There are conflicting studies on the effectiveness of hydrolyzed diets for allergic patients, with 10% to 40% of patients allergic to the basic protein continuing to show clinical signs on a hydrolyzed version⁷.

Home cooked diet is thus regarded as the most reliable diagnostic option, however less practical.

During the elimination diet, the dog must not receive any other food, treats, medications, vitamins, or supplements with any protein other than the one chosen for the diet.

Food elimination Trial - Food Challenge



Next+ Food Reactivity Test

Although not diagnostic, combined assays of food specific IgE and IgG, represent a useful tool to identify a suitable composition for an elimination diet, when attempting to diagnose and manage a suspected adverse food reaction¹⁰.



Which Elimination Diet should be used?

The testing of IgE on its own, represents a half-reliable tool. Evidence shows significant improvements when choosing an elimination diet based on combination of IgE and IgG results, when compared with IgE alone^{10,11}.

Generally, serum testing for food-specific IgE and IgG shown an average accuracy of 75%¹². High predictive negative values (Ø75% IgE and 84% IgG) represent a more accurate interpretation when compared with positive predictability (15-100% IgE and 35% IgG)¹².

Next+ Food Reactivity Test includes the results for food-specific IgE and IgG levels for 20 different allergens, together with clear advice on potential suitable ingredients and commercial diets for a food elimination trial.



23 allergens x2 (IgE + IgG)

- Beef
- Chicken
- Lamb
- Pork
- Turkey
- Venison
- Horse
- Duck
- Rabbit
- White fish
- Blue fish
- Milk
- Egg
- Soya bean
- Corn
- Potato
- Sugar beet
- Carrot
- Pea
- Wheat
- Rice
- Oat
- Yeast




Management

Once food allergy is established, strict avoidance of allergen is the gold standard. Taxonomic relationship between protein sources should be considered to avoid the risk of cross-reactivity¹³.

If the trial appears to have been performed correctly, but the patient did not improve or slightly improved, then it is likely suffering from atopic dermatitis and reacting to environmental allergens. Next+ Serum Test and/or Artuvetrin® Skin Test can identify these

environmental allergens that your patient is reacting to and help you putting together an appropriate immunotherapy to treat the cause of the allergy – Artuvetrin® Therapy.



References

- 1 Sampson, H. A., 2004: Update on food allergy. *Journal of Allergy Clinical Immunology*. 113:805-819
- 2 Verlinden, A. et al, 2006: Food allergy in dogs and cats: a review. *Clinical Reviews in Food Science and Nutrition*. 46:259-273
- 3 Olivry, T., Mueller, R., 2017: Critically appraised topic on adverse food reactions of companion animals (3): prevalence of cutaneous adverse food reactions in dogs and cats. *BMC Veterinary Research*. 13:51
- 4 Pali-Schöll, I. et al, 2017: Comparing immediate-type food allergy in humans and companion animals. *EAACI*. 72:1643-1656
- 5 Jeffers J. G. et al, 1996: Responses of dogs with food allergies to single-ingredient dietary provocation. *J Am Vet Med Assoc*. 209:608-611
- 6 Ricci, R., et al, 2010: A comparison of the clinical manifestations of feeding whole and hydrolyzed chicken to dogs with hypersensitivity to the native protein. *Vet Dermatol*. 21:358-366
- 7 Olivry, T. et al, 2010: A systematic review of the evidence of reduced allergenicity and clinical benefit of food hydrolysates in dogs with cutaneous adverse food reactions. *Vet Dermatol*. 21:32-41
- 8 Ricci, R. et al, 2013: Identification of undeclared sources of animal origin in canina dry foods used in dietary elimination trial. *J Anim Physiol Anim Nutr (Berl)*. 97(Suppl 1):32-38
- 9 Raditic, D. et al, 2011: ELISA testing for common food antigens in four dry foods used in dietary elimination trial. *J Anim Physiol Anim Nutr (Berl)*. 95:90-97
- 10 Haliwell, R. et al, 2004: IgE and IgG antibodies to food antigens in sera from normal dogs, atopic dogs and dogs with adverse food reactions. *Vet Dermatol*. 15: 2-3
- 11 Arribas, J. et al: Assessment of an "in-vitro" diagnosis of IgG/IgE in dogs suspected to suffer food allergy and/or food intolerance. School of Veterinary Science Madrid.
- 12 Olivry, T., Mueller, R., 2017: Critically appraised topic on adverse food reactions of companion animals (4): can we diagnose adverse food reactions in dogs and cats with in vivo or in vitro tests? *BMC Veterinary Research*. 13:275
- 13 Bexley, J. et al, 2016: Serological cross reactivity between beef, lamb and cow's milk allergen extracts in dogs. *Vet Dermatol*. 27:9



nextmune

artuvet | alergovet | dr baddaky

